

Three-Loop HTL Free Energy for QED

Nan Su

Frankfurt Institute for Advanced Studies, Goethe-Universität Frankfurt

Collaborators: Michael Strickland (Gettysburg and FIAS), Jens Andersen (NTNU)

Reference: arXiv:0906.2936

The 5-th International Conference on Quarks and Nuclear Physics
Beijing, September 24, 2009




FIAS

Frankfurt Institute for Advanced Studies

Contents

- Introduction
 - Poor convergence of naive perturbation theory at finite temperature
- Hard-Thermal-Loop Perturbation Theory
- Conclusions and Outlook

Introduction



Perturbative QCD free energy with $N_c = 3$ and $N_f = 2$ vs temperature. ($\pi T \leq \mu \leq 4\pi T$)
4-d lattice results from Karsch et al, 03.

(Here $\alpha_s = g_s^2/4\pi$)

- The weak-coupling expansion of the QCD free energy, \mathcal{F} , has been calculated to order $\alpha_s^3 \log \alpha_s$.^{1,2,3,4}
- At temperatures expected at RHIC energies, $T \sim 0.3$ GeV, the running coupling constant $\alpha_s(2\pi T)$ is approximately 1/3, or $g_s \sim 2$.
- The successive terms contributing to \mathcal{F} can strictly only form a decreasing series if $\alpha_s \lesssim 1/20$ which corresponds to $T \sim 10^5$ GeV.


¹ Arnold and Zhai, 94/95.

² Kastening and Zhai, 95.

³ Braaten and Nieto, 96.

⁴ Kajantie, Laine, Rummukainen and Schröder, 02.

Introduction




LO and NLO HTLpt free energy of QCD with $N_c = 3$ and $N_f = 2$
together with the perturbative prediction accurate to g^5 .

- Hard-thermal-loop (HTL) perturbation theory ^{4,5} is a systematic, self-consistent and gauge-invariant reorganization of thermal quantum fields.
- Hard-thermal-loop perturbation theory is formulated in Minkowski space, therefore it is in principle possible to carry out real time calculations.
- Interested in $T > 2 - 3 T_c$.

⁴ Andersen, Braaten, Strickland, 99/99/99.

⁵ Andersen, Braaten, Petitgirard, Strickland, 02;
Andersen, Petitgirard, Strickland, 03.

But there is still work to do!



LO and NLO HTLpt free energy of QCD with $N_c = 3$ and $N_f = 2$
together with the perturbative prediction accurate to g^5 .

- g^4 and g^5 terms can't be fully fixed at NLO. Some of them enter at NNLO. The result has the right magnitude, but the wrong sign.
- Running coupling effect doesn't enter at NLO. At this order, running coupling needs to be put by hand. Coupling constant renormalization enters at NNLO as well.

NNLO is needed!

Hard-Thermal-Loop Perturbation Theory (HTLpt)

- Hard-thermal-loop perturbation theory is a reorganization of the perturbative series for QCD

$$\mathcal{L}_{\text{HTLpt}} = (\mathcal{L}_{\text{QCD}} + \mathcal{L}_{\text{HTL}}) \Big|_{g \rightarrow \sqrt{\delta} g} + \Delta \mathcal{L}_{\text{HTL}}$$

The HTL “improvement” term is


$$\mathcal{L}_{\text{HTL}} = -\frac{1}{2}(1 - \delta)m_D^2 \text{Tr} \left(G_{\mu\alpha} \left\langle \frac{y^\alpha y^\beta}{(y \cdot D)^2} \right\rangle_y G^\mu{}_\beta \right)$$

where $\langle \dots \rangle_y$ indicates angle average

HTLpt: 1-loop free energy for pure glue

- Separation into hard and soft contributions ($d = 3 - 2\epsilon$)

$$\mathcal{F}_g = -\frac{1}{2} \oint_P \{ (d-1) \log[-\Delta_T(P)] + \log \Delta_L(P) \}$$



- Hard momenta ($\omega, \mathbf{p} \sim T$)

$$\begin{aligned} \mathcal{F}_g^{(h)} = & \frac{d-1}{2} \oint_P \log(P^2) + \frac{1}{2} m_D^2 \oint_P \frac{1}{P^2} - \frac{1}{4(d-1)} m_D^4 \oint_P \left[\frac{1}{(P^2)^2} \right. \\ & \left. - 2 \frac{1}{p^2 P^2} - 2d \frac{1}{p^4} \mathcal{T}_P + 2 \frac{1}{p^2 P^2} \mathcal{T}_P + d \frac{1}{p^4} (\mathcal{T}_P)^2 \right] + \mathcal{O}(m_D^6) \end{aligned}$$

- Soft momenta ($\omega, \mathbf{p} \sim gT$)

$$\mathcal{F}_g^{(s)} = \frac{1}{2} T \int_{\mathbf{p}} \log(p^2 + m_D^2)$$

HTLpt: 1-loop free energy for pure glue

- LO thermodynamical potential


$$\frac{\Omega_{\text{LO}}}{\mathcal{F}_{\text{ideal}}} = 1 - \frac{15}{2}\hat{m}_D^2 + 30\hat{m}_D^3 + \frac{45}{4} \left(\log \frac{\hat{\mu}}{2} - \frac{7}{2} + \gamma + \frac{\pi^2}{3} \right) \hat{m}_D^4 + \mathcal{O}(\hat{m}_D^6),$$

where $\hat{m}_D = \frac{m_D}{2\pi T}$ and $\hat{\mu} = \frac{\mu}{2\pi T}$.

- The gap equation is not well-defined at LO (α_s does not appear above). However, we can get LO free energy by setting


$$m_D = gT.$$

HTLpt: 1- and 2-loop free energy for pure glue




LO and NLO HTLpt free energy of pure glue vs temperature
Andersen, Braaten, Petitgirard, Strickland, 02.

HTLpt: naive pert. expansion of QED free energy



Perturbative QED free energy (Kastening and Zhai, 95)

HTLpt: 3-loop diagrams for QED



3-loop QED diagrams contributing to HTLpt

HTLpt: Counterterms

- The counterterms we need in the 3rd loop renormalization are

$$\Delta\mathcal{E}_0 = \frac{1}{128\pi^2\epsilon} m_D^4$$

$$\Delta m_D^2 = -N_f \frac{\alpha}{3\pi\epsilon} m_D^2$$

$$\Delta m_f^2 = \frac{3\alpha}{4\pi\epsilon} m_f^2$$

$$\Delta\alpha = N_f \frac{\alpha^2}{3\pi\epsilon} \text{ (same as zero T!)}$$

HTLpt: 3-loop thermodynamic potential for QED

- The NNLO thermodynamic potential reads

$$\begin{aligned}\Omega_{\text{NNLO}} = & -\frac{\pi^2 T^4}{45} \left\{ 1 + \frac{7}{4} N_f - \frac{15}{4} \hat{m}_D^3 \right. \\ & + N_f \frac{\alpha}{\pi} \left[-\frac{25}{8} + \frac{15}{2} \hat{m}_D + 15 \left(\log \frac{\hat{\mu}}{2} - \frac{1}{2} + \gamma + 2 \log 2 \right) \hat{m}_D^3 - 90 \hat{m}_D \hat{m}_f^2 \right] \\ & + N_f \left(\frac{\alpha}{\pi} \right)^2 \left[\frac{15}{64} (35 - 32 \log 2) - \frac{45}{2} \hat{m}_D \right] \\ & + N_f^2 \left(\frac{\alpha}{\pi} \right)^2 \left[\frac{25}{12} \left(\log \frac{\hat{\mu}}{2} + \frac{1}{20} + \frac{3}{5} \gamma - \frac{66}{25} \log 2 + \frac{4}{5} \frac{\zeta'(-1)}{\zeta(-1)} - \frac{2}{5} \frac{\zeta'(-3)}{\zeta(-3)} \right) \right. \\ & \left. \left. + \frac{5}{4} \frac{1}{\hat{m}_D} - 15 \left(\log \frac{\hat{\mu}}{2} - \frac{1}{2} + \gamma + 2 \log 2 \right) \hat{m}_D + 30 \frac{\hat{m}_f^2}{\hat{m}_D} \right] \right\}\end{aligned}$$


PURELY ANALYTIC!

- To eliminate the m_D and m_f dependence, the gap equations are imposed

$$\frac{\partial}{\partial m_D} \Omega(T, \alpha, m_D, m_f, \delta = 1) = 0$$


$$\frac{\partial}{\partial m_f} \Omega(T, \alpha, m_D, m_f, \delta = 1) = 0$$

HTLpt: 2- and 3-loop free energy for QED



NLO and NNLO HTLpt predictions for QED free energy

HTLpt: comparison of different schemes



Comparison of three different predictions for QED free energy at $\mu = 2\pi T$


3-loop Φ -derivable result is taken from Andersen and Strickland, 05

Conclusions and Outlook

- The problem of bad convergence of weak-coupling expansion at finite temperature is generic.
- It does not just happen in gauge theories, but also in scalar theories, and even in quantum mechanics.
- Hard-thermal-loop perturbation theory, which is formulated in Minkowski space, can improve the convergence of perturbative calculations in a gauge-invariant manner.
- By pushing forward, hopefully, hard-thermal-loop perturbation theory can provide a generic way towards a convergent gauge theory at high temperature, $T > 2 - 3 T_c$.
- Once the NNLO QCD thermodynamics is obtained, we can begin to calculate dynamic quantities.


Back-up

Large-N Limit



QED free energy in the large N_f limit. The exact result is taken from Ipp, Moore and Rebhan 03.

Screened Perturbation Theory




4-loop SPT pressure vs weak-coupling pressure

Andersen, Braaten and Strickland, 00. Andersen and Strickland, 01.


Andersen and Kyllingstad, 08.

HTLpt: 3-loop free energy for QED



NLO and NNLO HTLpt thermodynamic potentials with perturbative thermal masses

HTLpt: 3-loop free energy for QED



The imaginary part of NNLO HTLpt predictions for QED free energy

Weinberg and Wu, 87